



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### (Chloromethyl)trimethylsilane at 160 K

**Citation for published version:**

Fraser, S & Parsons, S 2004, '(Chloromethyl)trimethylsilane at 160 K' Acta Crystallographica Section E: Structure Reports Online, vol. 60, pp. O564-O565. DOI: 10.1107/S1600536804005276

**Digital Object Identifier (DOI):**

[10.1107/S1600536804005276](https://doi.org/10.1107/S1600536804005276)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Acta Crystallographica Section E: Structure Reports Online

**Publisher Rights Statement:**

Copyright © 2004 International Union of Crystallography; all rights reserved.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## (Chloromethyl)trimethylsilane at 160 K

Stewart Fraser and Simon Parsons\*

School of Chemistry, The University of Edinburgh, King's Buildings, West Mains Road, Edinburgh EH9 3JJ, Scotland

Correspondence e-mail: s.parsons@ed.ac.uk

## Key indicators

Single-crystal X-ray study

$T = 160\text{ K}$

Mean  $\sigma(\text{Si}-\text{C}) = 0.004\text{ \AA}$

$R$  factor = 0.040

$wR$  factor = 0.115


Data-to-parameter ratio = 23.3

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

(Chloromethyl)trimethylsilane,  $\text{Me}_3\text{SiCH}_2\text{Cl}$  or  $\text{C}_4\text{H}_{11}\text{ClSi}$ , is a liquid at room temperature, and it was crystallized using *in situ* methods. The  $\text{C}-\text{Si}-\text{C}$  bond angles involving the chloromethyl group are somewhat smaller than those involving only methyl groups [ $105.5(2)$ – $109.47(19)^\circ$  versus  $110.01(19)$ – $111.2(2)^\circ$ ], which is ascribable to both the electronegative and the steric effects of the Cl atom.

## Comment

(Chloromethyl)trimethylsilane, (I), is a liquid under ambient conditions, and a crystal was obtained by *in situ* crystallization of a sample held in a hand-drawn Pyrex capillary (Boese & Nussbaumer, 1994).



Molecules of (I) adopt the expected tetrahedral configuration at Si (Fig. 1).  $\text{Si}-\text{C}$  bond distances fall into the range  $1.848(4)$ – $1.880(4)\text{ \AA}$ , although to within experimental error the bond distances and angles have  $C_s$  symmetry, with a mirror plane passing through atoms Si1, C1, Cl1 and C4; the  $\text{C4}-\text{Si1}-\text{C1}-\text{Cl1}$  torsion angle [ $175.3(2)^\circ$ ] shows a somewhat more significant deviation from the symmetry. The bond angles at atom Si1 involving the more electronegative  $\text{CH}_2\text{Cl}$  group are smaller [ $105.5(2)$ – $109.47(19)^\circ$ ] than those involving only methyl groups [ $110.01(19)$ – $111.2(2)^\circ$ ]. The smaller




Figure 1

A view of the molecular structure of (I). Displacement ellipsoids are drawn at the 50% probability level and H atoms are represented as spheres of arbitrary radii.

Received 16 February 2004

Accepted 8 March 2004

Online 24 March 2004

magnitude of C1—Si1—C4 [105.5 (2)°] relative to C1—Si1—C2 [109.4 (2)°] and C1—Si1—C3 [109.47 (19)°] presumably reflects the steric influence of the Cl atom.

The only intermolecular interactions falling within the sum of the van der Waals radii (Bondi, 1964) of the participating atoms are weak Cl1...H33<sup>i</sup> [symmetry code (i):  $-x, -y, z - \frac{1}{2}$ ] interactions (2.93 Å; the sum of the van der Waals radii of Cl and H is 2.95 Å). These result in chains that spiral about the 2<sub>1</sub> axis parallel to the *c* direction (Fig. 2).

## Experimental

A sample of (I) was obtained from Aldrich and used as received. Compound (I) is a liquid under ambient conditions and it was crystallized *in situ* in a capillary (o.d. 0.34 mm) mounted on the diffractometer. A crystal was grown by first establishing a seed in a small volume of the liquid at 182.8 K, and then cooling at a rate of 10 K h<sup>-1</sup>. The sample was then cooled to 160 K for data collection.

### Crystal data

C <sub>4</sub> H <sub>11</sub> ClSi	Mo K $\alpha$ radiation
$M_r = 122.67$	Cell parameters from 72 reflections
Orthorhombic, <i>Pna</i> 2 <sub>1</sub>	$\theta = 15\text{--}16^\circ$
$a = 13.8776$ (13) Å	$\mu = 0.56\text{ mm}^{-1}$
$b = 6.3855$ (9) Å	$T = 160\text{ K}$
$c = 8.4000$ (10) Å	Cylinder, colourless
$V = 744.37$ (15) Å <sup>3</sup>	$0.50 \times 0.39 \times 0.39\text{ mm}$
$Z = 4$	$0.50\text{ mm length, }0.39\text{ mm radius}$
$D_x = 1.095\text{ Mg m}^{-3}$	

### Data collection

Stoe STADI-4 diffractometer equipped with an Oxford Cryo-systems low-temperature device (Cosier & Glazer, 1986)	1306 independent reflections
$\omega$ - $\theta$ scans	1186 reflections with $I > 2\sigma(I)$
Absorption correction: $\psi$ scan [azimuthal absorption correction (North <i>et al.</i> , 1968) applied using XPREP (Sheldrick, 1997)]	$R_{\text{int}} = 0.030$
$T_{\text{min}} = 0.741$ , $T_{\text{max}} = 0.804$	$\theta_{\text{max}} = 25.0^\circ$
3768 measured reflections	$h = -1 \rightarrow 16$
	$k = -7 \rightarrow 7$
	$l = -9 \rightarrow 9$
	3 standard reflections
	frequency: 60 min
	intensity decay: none


### Refinement

Refinement on $F^2$	where $P = 0.3333\max(0, F_o^2) + 0.6667F_c^2$
$R[F^2 > 2\sigma(F^2)] = 0.040$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$wR(F^2) = 0.115$	$\Delta\rho_{\text{max}} = 0.43\text{ e \AA}^{-3}$
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.32\text{ e \AA}^{-3}$
1305 reflections	Absolute structure: Flack (1983),
56 parameters	602 Friedel pairs
H-atom parameters constrained	Flack parameter = $-0.17$ (17)
$w = 1/[\sigma^2(F^2) + (0.0706P)^2 + 0.41P]$	

**Table 1**

Selected geometric parameters (Å, °).

Cl1—C1	1.798 (5)	Si1—C3	1.862 (4)
Si1—C1	1.880 (4)	Si1—C4	1.867 (4)
Si1—C2	1.848 (4)		
C1—Si1—C2	109.4 (2)	C2—Si1—C4	111.2 (2)
C1—Si1—C3	109.47 (19)	C3—Si1—C4	111.18 (19)
C1—Si1—C4	105.5 (2)	Cl1—C1—Si1	111.9 (2)
C2—Si1—C3	110.01 (19)		



**Figure 2**

The molecular packing of (I), viewed along the *b* axis. Weak intermolecular Cl1...H33 interactions are shown as dotted lines.

The positions of the H atoms were recalculated geometrically after each refinement cycle, using a C—H distance of 1.00 Å, and they were assigned  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The 10,1,0 reflection was omitted as an outlier.

Data collection: *DIF4* (Stoe & Cie, 1990); cell refinement: *DIF4*; data reduction: *REDU4* (Stoe & Cie, 1990); program(s) used to solve structure: *DIRDIF* (Beurskens *et al.*, 1996); program(s) used to refine structure: *CRYSTALS* (Betteridge *et al.*, 2003); molecular graphics: *CAMERON* (Watkin *et al.*, 1996) and *XP* (Sheldrick, 1997); software used to prepare material for publication: *CRYSTALS*, *enCIFer* (CCDC, 2003) and *PLATON* (Spek, 2003) used within *WinGX* (Farrugia, 1999).

The authors thank the EPSRC and the University of Edinburgh for funding.

## References

- Betteridge, P. W., Carruthers, J. R., Cooper, R. I., Prout, C. K. & Watkin, D. J. (2003). *J. Appl. Cryst.* **36**, 1487.
- Beurskens, P. T., Admiraal, G., Beurskens, G., Bosman, W. P., García-Granda, S., Gould, R. O., Smits, J. M. M. & Smykalla, C. (1996). *The DIRDIF96 Program System*. Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
- Boese, R. & Nussbaumer, M. (1994). *Correlations, Transformations and Interactions in Organic Crystal Chemistry, IUCr Crystallographic Symposia*, Vol. 7, edited by D. W. Jones & A. Katrusiak, pp. 20–37. Oxford University Press.
- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
- CCDC (2003). *enCIFer*. Test version. Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, England.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (1997). *XP, XPREP and SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Stoe & Cie (1990). *DIF4* (Version 7.09/DOS) and *REDU4* (Version 7.03/DOS). Stoe & Cie, Darmstadt, Germany.
- Watkin, D. J., Prout, C. K. & Pearce, L. J. (1996). *CAMERON*. Chemical Crystallography Laboratory, University of Oxford, England.